arXiv:0910.4124v2 [math.DG] 19 Feb 2010

MINIMAL SURFACES IN $\mathbb{R}^3$ PROPERLY PROJECTING INTO $\mathbb{R}^2$

ANTONIO ALARCÓN AND FRANCISCO J. LÓPEZ

ABSTRACT. For all open Riemann surface $\mathcal{N}$ and real number $\theta \in (0, \pi/4)$, we construct a conformal minimal immersion $X = (X_1, X_2, X_3) : \mathcal{N} \to \mathbb{R}^3$ such that $X_3 + \tan(\theta)|X_1| : \mathcal{N} \to \mathbb{R}$ is positive and proper. Furthermore, X can be chosen with arbitrarily prescribed flux map.

Moreover, we produce properly immersed hyperbolic minimal surfaces with non empty boundary in $\mathbb{R}^3$ lying above a negative sublinear graph.

1. INTRODUCTION

The conformal structure of a complete minimal surface plays a fundamental role in its global properties. It is then important to determine the conformal type of a given minimal surface. An open Riemann surface is said to be *hyperbolic* if an only if it carries a negative non-constant subharmonic function. Otherwise, it is said to be *parabolic*. Compact Riemann surfaces with empty boundary are said to be *elliptic*.

Complete minimal surfaces with finite total curvature or complete embedded minimal surfaces with finite topology in $\mathbb{R}^3$ are properly immersed and have parabolic conformal type (for further information, see [**Os**, **JM**, **CM**, **MPR**, **MP2**]). On the other hand, there exists properly immersed hyperbolic minimal surfaces in $\mathbb{R}^3$ with arbitrary non compact topology [**Mo**, **AFM**, **FMM**].

It is then interesting to elucidate how properness and completeness influence the conformal geometry of minimal surfaces. In [**Lo2**] it is shown that any open Riemann surface admits a conformal complete minimal immersion in $\mathbb{R}^3$, even with arbitrarily prescribed flux map. In this paper we extend this result to the family of proper minimal immersions, proving considerably more (see Theorem 4.4):

> **Theorem I.** *For all open Riemann surface $\mathcal{N}$, group morphism $p : \mathcal{H}_1(\mathcal{N}, \mathbb{Z}) \to \mathbb{R}^3$ and real number $\theta \in (0, \frac{\pi}{4})$, there exists a conformal minimal immersion $X = (X_1, X_2, X_3) :$ $\mathcal{N} \to \mathbb{R}^3$ satisfying that:*
> - *$X_3 + \tan(\theta)|X_1| : \mathcal{N} \to \mathbb{R}$ is positive and proper, and*
> - *$\int_\gamma \partial X = ip(\gamma)$ for all $\gamma \in \mathcal{H}_1(\mathcal{N}, \mathbb{Z})$, where ∂ is the complex differential operator.*

This result is sharp in the sense that the angle θ cannot be zero. Indeed, by the Strong Half Space Theorem [**HM**] properly immersed minimal surfaces in a half space are planes. Contrariwise, Theorem I shows that any wedge of angle greater than π in $\mathbb{R}^3$ contains minimal surfaces properly immersed in $\mathbb{R}^3$, even of hyperbolic type. In particular, neither open wedges nor closed wedges of angle greater than π are universal regions for surfaces (see [**MP1**] for a good setting). Other Picard conditions for properly immersed minimal surfaces in $\mathbb{R}^3$ guaranteeing parabolicity can be found in [**Lo1**].

2000 *Mathematics Subject Classification.* 53A10; 49Q05, 49Q10, 53C42.

Key words and phrases. Proper minimal surfaces, Riemann surfaces of arbitrary conformal structure.

Research of both authors is partially supported by MCYT-FEDER research project MTM2007-61775 and Junta de Andalucía Grant P09-FQM-5088.

From Theorem I follow some remarkable results concerning not only minimal surfaces. We are going to mention three of them related to proper harmonic maps into $\mathbb{C}$, proper holomorphic null curves in $\mathbb{C}^3$ and maximal surfaces in the Lorentz-Minkowski space $\mathbb{R}^3_1$.

Schoen and Yau conjectured that there are no proper harmonic maps from $\mathbb{D}$ to $\mathbb{C}$ with flat metrics, and connected this question with the existence of hyperbolic minimal surfaces in $\mathbb{R}^3$ properly projecting into $\mathbb{R}^2$ [**SY**, p. 18]. A counterexample to this conjecture follows from the results in [**DF**], which imply the existence of proper harmonic maps from any finite bordered Riemann surface (that is to say, a compact Riemann surface minus a finite collection of pairwise disjoint closed discs) into $\mathbb{R}^2$. It remains open whether or not a hyperbolic minimal surface in $\mathbb{R}^3$ can be properly projected into $\mathbb{R}^2$. Theorem I answers positively this question for minimal surfaces with arbitrary open conformal structure (just notice that the harmonic map $(X_1, X_3) : \mathcal{N} \to \mathbb{R}^2$ is proper).

It is well known that any open Riemann surface properly holomorphically embeds in $\mathbb{C}^3$ and immerses in $\mathbb{C}^2$ [**Bi**, **Nar**, **Re**]. Moreover, there are proper null immersions in $\mathbb{C}^3$ of the unit disc [**Mo**], and of any open parabolic Riemann surface of finite topology [**Pi**, **Lo2**]. Theorem I also shows that any open Riemann surface admits a proper *null* immersion in $\mathbb{C}^3$, and a holomorphic immersion in $\mathbb{C}^2$ properly projecting into $\mathbb{R}^2$. Indeed, choosing $p = 0$ in Theorem I and labeling $X^* = (X_1^*, X_2^*, X_3^*)$ as the conjugate minimal immersion of X, the map $X + iX^* = (F_1, F_2, F_3) : \mathcal{N} \to \mathbb{C}^3$ is a proper holomorphic null immersion, and $(F_1, F_3) : \mathcal{N} \to \mathbb{C}^2$ is a holomorphic immersion which properly projects into $\mathbb{R}^2$.

Finally, from Theorem I follows the existence of proper Lorentzian null holomorphic immersions in $\mathbb{C}^3$ (see [**UY**]) and proper conformal maximal immersions in the Lorentz-Minkowski space, with singularities and arbitrary conformal structure. See [**Al**] for the hyperbolic simply connected case.

The last part of the paper is devoted to properly immersed minimal surfaces in $\mathbb{R}^3$ with non empty boundary. A Riemann surface M with *non empty boundary* is said to be *parabolic* if bounded harmonic functions on M are determined by their boundary values, or equivalently, if the harmonic measure of M respect to a point $P \in M - \partial(M)$ is full on $\partial(M)$. Otherwise, the surface is said to be *hyperbolic* (see [**AS**, **Pe**] for a good setting). For instance, $\overline{\mathbb{D}} - \{1\}$ is parabolic whereas $\overline{\mathbb{D}}_+ := \overline{\mathbb{D}} \cap \{z \in \mathbb{C} \,|\, \mathrm{Im}(z) > 0\}$ is hyperbolic. Properly immersed minimal surfaces with non empty boundary lying in a half space of $\mathbb{R}^3$ are parabolic [**CKMR**], and the same result holds for proper minimal graphs in $\mathbb{R}^3$ [**Ne**]. It is also known that any properly immersed minimal surface in $\mathbb{R}^3$ with non empty boundary lying over a negative sublinear graph in $\mathbb{R}^3$ and whose Gaussian image is contained in a hyperbolic domain of the Riemann sphere is parabolic [**LP**]. We prove the following complementary result (see Theorem 5.1), which also shows that the condition about the size of the Gauss map in [**LP**] plays an important role:

> **Theorem II.** *There exists a conformal minimal immersion* $X = (X_1, X_2, X_3) : \overline{\mathbb{D}}_+ \to \mathbb{R}^3$ *such that* $(X_1, X_3) : \overline{\mathbb{D}}_+ \to \mathbb{R}^2$ *is proper and* $\lim_{n\to\infty} \min\{\frac{X_3(p_n)}{|X_1(p_n)|+1}, 0\} = 0$ *for all divergent sequence* $\{p_n\}_{n\in\mathbb{N}}$ *in* $\overline{\mathbb{D}}_+$.

Theorem II contributes to the understanding of Meeks' conjecture about parabolicity of minimal surfaces with boundary. This conjecture asserts that any properly immersed minimal surface lying above a negative half catenoid is parabolic.

The techniques developed in this paper may be applied to a wide range of problems on minimal surfaces theory. In the paper [**AFL**] complete minimal surfaces in $\mathbb{R}^3$ with arbitrary conformal structure and whose Gauss map misses two points are constructed. Our tools come from deep results on approximation theory by meromorphic functions [**Sc1**, **Sc2**, **Ro**]. The most useful one is the Approximation Lemma in Section 2, where accurate use of Runge's approximation theorem and classical theory of Riemann surfaces [**AS**, **FK**] is made. In this way, we can refine the classical construction methods of complete minimal surfaces (see, among others, **[JX**, **Nad**, **LMM**, **MUY]** for a good setting).

The paper is laid out as follows. In Section 2 we introduce the necessary background on Riemann surfaces and approximation theory, and the required notations. Furthermore, we prove the Approximation Lemma. Section 3 is devoted to some preliminaries on minimal surfaces in $\mathbb{R}^3$. Finally, Theorems I and II are proved in Sections 4 and 5, respectively.

2. RIEMANN SURFACES AND APPROXIMATION RESULTS

Given a topological surface W, $\partial(W)$ will denote the one dimensional topological manifold determined by the boundary points of W. Given $S \subset W$, call by S° and $\overline{S}$ the interior and the closure of S in W, respectively.

A Riemann surface M is said to be *open* if it is non-compact and $\partial(M) = \emptyset$. As usual, $\overline{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ will denote the Riemann sphere. We denote ∂ as the global complex operator given by $\partial|_U = \frac{\partial}{\partial z} dz$ for any conformal chart (U, z) on M.

Remark 2.1. Throughout this paper $\mathcal{N}$ will denote a fixed but arbitrary open Riemann surface.

Let S denote a subset of $\mathcal{N}$, $S \neq \mathcal{N}$. We denote by $\mathcal{F}_0(S)$, respectively $\mathcal{F}(S)$, as the space of continuous functions $f : S \to \mathbb{C}$, respectively $f : S \to \overline{\mathbb{C}}$, which are holomorphic, respectively meromorphic, on an open neighborhood of S in $\mathcal{N}$, and $f^{-1}(\infty) \subset S^\circ$. Likewise, $\mathcal{F}_0^*(S)$, respectively $\mathcal{F}^*(S)$, will denote the space of continuous functions $f : S \to \mathbb{C}$, respectively $f : S \to \overline{\mathbb{C}}$, being holomorphic, respectively meromorphic, on S° and satisfying that $f^{-1}(\infty) \subset S^\circ$.

As usual, a 1-form θ on S is said to be of type $(1,0)$ if for any conformal chart (U, z) in $\mathcal{N}$, $\theta|_{U \cap S} = h(z)dz$ for some function $h : U \cap S \to \overline{\mathbb{C}}$. We denote by $\Omega_0(S)$, respectively $\Omega(S)$, as the space of holomorphic, respectively meromorphic, 1-forms on an open neighborhood of S in $\mathcal{N}$, and without poles on $\overline{S} - S^\circ$. We call $\Omega^*(S)$ as the space of 1-forms θ of type $(1,0)$ on S such that $(\theta|_U)/dz \in \mathcal{F}^*(S \cap U)$ for any conformal chart (U, z) on $\mathcal{N}$. Likewise we define $\Omega_0^*(S)$.

Let $\mathfrak{Div}(S)$ denote the free commutative group of divisors of S with multiplicative notation. If $D = \prod_{i=1}^n Q_i^{n_i} \in \mathfrak{Div}(S)$, where $n_i \in \mathbb{Z} - \{0\}$ for all i, the set $\{Q_1, \ldots, Q_n\}$ is said to be the support of D, written $\operatorname{supp}(D)$. A divisor $D \in \mathfrak{Div}(S)$ is said to be *integral* if $D = \prod_{i=1}^n Q_i^{n_i}$ and $n_i \geq 0$ for all i. Given D_1, $D_2 \in \mathfrak{Div}(S)$, $D_1 \geq D_2$ if and only if $D_1 D_2^{-1}$ is integral. For any $f \in \mathcal{F}(S)$ we denote by $(f)_0$ and $(f)_\infty$ its associated integral divisors of zeroes and poles in S, respectively, and label $(f) = \frac{(f)_0}{(f)_\infty}$ as the divisor associated to f on S. Likewise we define $(\theta)_0$, $(\theta)_\infty$ for any $\theta \in \Omega(S)$, and call $(\theta) = \frac{(\theta)_0}{(\theta)_\infty}$ as the divisor of θ on S.

In the sequel we will assume that S is a *compact subset* of $\mathcal{N}$ and $W \subset \mathcal{N}$ an open subset containing S.

By definition, a connected component V of $W - S$ is said to be *bounded* in W if $\overline{V} \cap W$ is compact, where $\overline{V}$ is the closure of V in $\mathcal{N}$.

Definition 2.2. A compact subset $S \subset W$ is said to be admissible in W if and only if:

- $W - S$ has no bounded components in W,
- $M_S := \overline{S^\circ}$ consists of a finite collection of pairwise disjoint compact regions in W with $\mathcal{C}^0$ boundary,
- $C_S := \overline{S - M_S}$ consists of a finite collection of pairwise disjoint analytical Jordan arcs, and
- any component α of C_S with an endpoint $P \in M_S$ admits an analytical extension β in W such that the unique component of $\beta - \alpha$ with endpoint P lies in M_S.

Observe that if S is admissible in $\mathcal{N}$ then it is admissible in W as well, but the contrary is in general false.

We shall say that a function $f \in \mathcal{F}^*(S)$, respectively $f \in \mathcal{F}_0^*(S)$, can be uniformly approximated on S by functions in $\mathcal{F}(W)$, respectively in $\mathcal{F}_0(W)$, if there exists $\{f_n\}_{n\in\mathbb{N}} \subset \mathcal{F}(W)$, respectively $\{f_n\}_{n\in\mathbb{N}} \subset \mathcal{F}_0(W)$, such that $\{|f_n - f|\}_{n\in\mathbb{N}} \to 0$ uniformly on S. We also say that $\{f_n|_S\}_{n\in\mathbb{N}} \to f$ in the *ω-topology*. In particular all f_n have the same set $\mathcal{P}_f$ of poles on S°. A 1-form $\theta \in \Omega^*(S)$, respectively $\theta \in \Omega_0^*(S)$, can be uniformly approximated on S by 1-forms in $\Omega(W)$, respectively in $\Omega_0(W)$, if there exists $\{\theta_n\}_{n\in\mathbb{N}} \subset \Omega(W)$, respectively $\{\theta_n\}_{n\in\mathbb{N}} \subset \Omega_0(W)$, such that $\{\frac{\theta_n-\theta}{dz}\}_{n\in\mathbb{N}} \to 0$ uniformly on $S \cap U$, for any conformal closed disc (U, dz) on W. In particular all θ_n have the same set of poles $\mathcal{P}_\theta$ on S°. As above, we say that $\{\theta_n|_S\}_{n\in\mathbb{N}} \to \theta$ in the ω-topology.

Recall that a compact Jordan arc in $\mathcal{N}$ is said to be analytical if it is contained in an open analytical Jordan arc in $\mathcal{N}$.

Given an admissible compact set $S \subset W$, a function $f : S \to \mathbb{C}^n$, $n \in \mathbb{N}$, is said to be smooth if $f|_{M_S}$ admits a smooth extension f_0 to an open domain V in W containing M_S, and for any component α of C_S and any open analytical Jordan arc β in W containing α, f admits an smooth extension f_β to β satisfying that $f_\beta|_{V\cap\beta} = f_0|_{V\cap\beta}$. A function $f \in \mathcal{F}^*(S)$ is said to be smooth if $f|_{S-V}$ is smooth, where V is any open neighbourhood of $f^{-1}(\infty)$ such that $S - V$ is admissible in $W - f^{-1}(\infty)$. Analogously, a 1-form $\theta \in \Omega^*(S)$ is said to be smooth if $(\theta|_{S\cap U})/dz \in \mathcal{F}^*(S \cap U)$ is smooth for any closed conformal disk (U, z) on W such that $S \cap U$ is an admissible set. Given a smooth $f \in \mathcal{F}^*(S)$, we set $df \in \Omega^*(S)$ as the smooth 1-form given by $df|_{M_S} = d(f|_{M_S})$ and $df|_{\alpha\cap U} = (f \circ \alpha)'(x)dz|_{\alpha\cap U}$, where $(U, z = x + iy)$ is a conformal chart on W such that $\alpha \cap U = z^{-1}(\mathbb{R} \cap z(U))$. Obviously, $df|_\alpha(t) = (f \circ \alpha)'(t)dt$ for any component α of C_S, where t is any smooth parameter along α.

A smooth 1-form $\theta \in \Omega^*(S)$ is said to be *exact* if $\theta = df$ for some smooth $f \in \mathcal{F}^*(S)$, or equivalently if $\int_\gamma \theta = 0$ for all $\gamma \in \mathcal{H}_1(S, \mathbb{Z})$.

Several extensions of classical Runge's Theorem can be found in [**Ro**, **Sc1**, **Sc2**]. For our purposes, we need only the following compilation result:

Theorem 2.3. *Let $S \subset W$ be a not necessarily connected admissible compact subset in W.*

Then any function $f \in \mathcal{F}^*(S)$ can be uniformly approximated on S by functions in $\mathcal{F}(W) \cap \mathcal{F}_0(W - \mathcal{P}_f)$, where $\mathcal{P}_f = f^{-1}(\infty)$. Furthermore, if $D \in \mathfrak{Div}(S)$ is an integral divisor satisfying that $supp(D) \subset S^\circ$, then the approximation sequence $\{f_n\}_{n\in\mathbb{N}}$ in $\mathcal{F}(W)$ can be chosen satisfying that $(f - f_n|_S)_0 \geq D$.

2.1. **The Approximation Lemmas.**

Throughout this section, $W \subset \mathcal{N}$ will denote an open connected subset of finite topology, and S an admissible compact subset in W.

Lemma 2.4. *Consider $f \in \mathcal{F}^*(S)$ such that f never vanishes on $S - S^\circ$. Then f can be uniformly approximated on S by functions $\{f_n\}_{n\in\mathbb{N}}$ in $\mathcal{F}(W)$ satisfying that $(f_n) = (f)$ on W for all n. In particular, f_n is holomorphic and never vanishing on $W - S$ for all n.*

Proof. Let μ and b denote the genus of W and the number of topological ends of $W - \text{supp}((f))$. It is well known (see [**FK**]) that there exist $2\mu + b - 1$ cohomologically independent 1-forms in $\Omega(W) \cap \Omega_0(W - \text{supp}((f)))$ generating the first holomorphic De Rham cohomology group $\mathcal{H}^1_{\text{hol}}(W - \text{supp}((f)))$. Furthermore, the 1-forms can be chosen having at most single poles at points of $\text{supp}((f))$. Thus, the map $H^1_{\text{hol}}(W - \text{supp}((f))) \longrightarrow \mathbb{C}^{2\mu+b-1}$, $\tau \mapsto \left(\int_c \tau\right)_{c\in B_0}$, where B_0 is any homology basis of $W - \text{supp}((f))$, is a linear isomorphism, and there exists $\tau \in \Omega(W) \cap$

$\Omega_0\big(W - \mathrm{supp}((f))\big)$ with at most single poles at points of $\mathrm{supp}((f))$ such that $\frac{1}{2\pi i}\int_\gamma \tau \in \mathbb{Z}$ for all $\gamma \in \mathcal{H}_1(W - \mathrm{supp}((f)), \mathbb{Z})$ and $df/f - \tau \in \Omega_0^*(S)$ is exact. Set $f_0 = f e^{-\int \tau} \in \mathcal{F}^*(S)$.

It is clear that $\log(f_0) \in \mathcal{F}_0^*(S)$, and so f_0 is holomorphic and never vanishing on S. By Theorem 2.3, there exists a sequence $\{h_n\}_{n\in\mathbb{N}} \subset \mathcal{F}_0^*(W)$ converging to $\log(f_0)$ in the ω-topology on S. The sequence $f_n = e^{h_n + \int \tau}$, $n \in \mathbb{N}$, solves the lemma. □

Lemma 2.5. *Consider $\theta \in \Omega^*(S)$ such that θ never vanishes on $S - S^\circ$.*

Then θ can be uniformly approximated on S by 1-forms $\{\theta_n\}_{n\in\mathbb{N}}$ in $\Omega(W)$ satisfying that $(\theta_n) = (\theta)$ on W. In particular, θ_n is holomorphic and never vanishing on $W - S$ for all n.

Proof. Let τ be a non zero 1-form in $\Omega_0(W)$ with finitely many zeroes, none of them lying in $S - S^\circ$. Label $f = \theta/\tau \in \mathcal{F}_*(S)$. By Lemma 2.4, f can be approximated in the ω-topology on S by a sequence $\{f_n\}_{n\in\mathbb{N}}$ in $\mathcal{F}(W)$ satisfying that $(f_n) = (f)$ on W for all n. It suffices to take $\theta_n := f_n\tau$, $n \in \mathbb{N}$. □

Lemma 2.6 (The Approximation Lemma)**.** *Let $\Phi = (\phi_j)_{j=1,2,3}$ be a smooth triple in $\Omega_0^*(S)^3$ such that $\sum_{j=1}^3 \phi_j^2 = 0$, $\sum_{j=1}^3 |\phi_j|^2$ never vanishes on S, and $\Phi|_{M_S} \in \Omega_0(M_S)^3$. Then Φ can be uniformly approximated on S by a sequence $\{\Phi_n = (\phi_{j,n})_{j=1,2,3}\}_{n\in\mathbb{N}} \subset \Omega_0(W)^3$ satisfying that:*

(i) $\sum_{j=1}^3 \phi_{j,n}^2 = 0$ and $\sum_{j=1}^3 |\phi_{j,n}|^2$ never vanishes on W,
(ii) $\Phi_n - \Phi$ is exact on S for all n.

Proof. Label $g = \frac{\phi_3}{\phi_1 - i\phi_2}$, $\eta_1 = \frac{1}{g}\phi_3 = \phi_1 - i\phi_2$ and $\eta_2 = g\phi_3 = -\phi_1 - i\phi_2 \in \Omega_0^*(S)$. We start with the following claim:

Claim 2.7. *Without loss of generality, we can assume that $g|_{M_S}$ is not constant.*

Proof. Suppose for a moment that $g|_{M_S}$ is constant, and up to replacing Φ by $\Phi \cdot A$ for a suitable orthogonal matrix $A \in \mathcal{O}(3,\mathbb{R})$, assume that $g \neq \infty$. For each $h \in \mathcal{F}_0(W)$, set $\eta_2(h) = (g+h)^2\eta_1$ and $\phi_3(h) = \eta_1(g+h)$. Let $\mathcal{B}$ be a homology basis of $\mathcal{H}_1(M_S, \mathbb{Z})$, label μ as its cardinal number and consider the holomorphic map $\mathcal{T} : \mathcal{F}_0(W) \to \mathbb{C}^{2\mu}$, $\mathcal{T}(h) = \big(\int_c (\eta_2(h) - \eta_2, \phi_3(h) - \phi_3)\big)_{c\in\mathcal{B}}$. Note that the analytical subset $\mathcal{T}^{-1}(0)$ is conical, that is to say, if $\mathcal{T}(h) = 0$ then $\mathcal{T}(\lambda h) = 0$ for all $\lambda \in \mathbb{C}$. Furthermore, since $\mathcal{F}_0(W)$ has infinite dimension we can choose a non constant $h \in \mathcal{T}^{-1}(0)$. Take $\{\lambda_n\}_{n\in\mathbb{N}} \subset \mathbb{C}$ converging to zero, set $h_n := \lambda_n h \in \mathcal{T}^{-1}(0)$ for all n, and notice that $\{h_n\}_{n\in\mathbb{N}} \to 0$ in the ω-topology on S.

Set $\Psi_n \equiv (\psi_{1,n}, \psi_{2,n}, \psi_{3,n}) := \big(\frac{1}{2}(\eta_1 - \eta_2(h_n)), \frac{i}{2}(\eta_1 + \eta_2(h_n)), \phi_3(h_n)\big) \in \Omega_0^*(S) \cap \Omega_0(M_S)^3$, and observe that $\sum_{j=1}^3 \psi_{j,n}^2 = 0$, $\sum_{j=1}^3 |\psi_{j,n}|^2$ never vanishes on S and $g_n = \frac{\psi_{3,n}}{\psi_{1,n} - i\psi_{2,n}}$ is holomorphic and non constant on M_S. It is clear that $\Psi_n - \Phi$ is exact on M_S, $n \in \mathbb{N}$. Furthermore, we can slightly deform $\Psi_n|_{C_S}$ so that $\Psi_n - \Phi$ is exact on S as well, $n \in \mathbb{N}$ (we leave the details to the reader). If the lemma holds for Ψ_n for all n, we can construct a sequence $\{\hat{\Psi}_{n,m}\}_{m\in\mathbb{N}} \subset \Omega_0^*(S)^3$ converging to Ψ_n in the ω-topology on S and satisfying that $\hat{\Psi}_{n,m} - \Psi_n$ is exact on S for all n. A standard diagonal argument proves the claim. □

Claim 2.8. *Without loss of generality, we can assume that g, $1/g$ and dg never vanish on $\partial(M_S) \cup C_S$ (hence the same holds for η_i, $i = 1,2$, and ϕ_j, $j = 1,2,3$). In particular, $g \in \mathcal{F}^*(S)$ and $dg \in \Omega^*(S)$.*

Proof. Take a sequence $M_1 \supset M_2 \supset \ldots$ of tubular neighborhoods of M_S in W such that $M_n \subset M_{n-1}^\circ$ for any n, $\cap_{n\in\mathbb{N}} M_n = M_S$, Φ (and so g) meromorphically extends (with the same name) to M_1, $\sum_{j=1}^3 |\phi_j|^2 \neq 0$ on M_1, and g, $1/g$, and dg never vanish on $\partial(M_n)$ for all n (take into account

Claim 2.7). Choose M_n in such a way that $S_n := M_n \cup C_S$ is an admissible subset of W and $\gamma - M_n^\circ$ is a (non-empty) Jordan arc for any component γ of C_S. In particular, $C_{S_n} = C_S - M_n^\circ$, $n \in \mathbb{N}$.

Let $(h_n, \psi_{3,n}) \in \mathcal{F}^*(S_n) \times \Omega_0^*(S_n)$ be smooth data such that

- $(h_n, \psi_{3,n})|_{M_{S_n}} = (g, \phi_3)|_{M_{S_n}}$ and $\sum_{j=1}^3 |\psi_{j,n}|^2$ never vanishes on S_n, where $\Psi_n = (\psi_{j,n})_{j=1,2,3} = \big(\frac{1}{2}(1/h_n - h_n), \frac{i}{2}(1/h_n + h_n), 1\big)\psi_{3,n} \in \Omega_0^*(S_n)^3$, $n \in \mathbb{N}$,
- h_n, $1/h_n$ and dh_n never vanish on $\partial(M_{S_n}) \cup C_{S_n}$,
- $\Psi_n|_S - \Phi$ is exact on S, and
- the sequence $\{\Psi_n|_S\}_{n\in\mathbb{N}} \subset \Omega_0^*(S)^3$ converges to Φ in the ω-topology on S.

The construction of these data is standard, we omit the details.

Label $\mathcal{T} \subset \Omega_0(W)^3$ as the subspace of data Ψ formally satisfying (i) and (ii). If the Lemma held for any of the data in $\{\Psi_n \mid n \in \mathbb{N}\}$, Ψ_n would lie in the closure of $\mathcal{T}$ in $\Omega_0^*(S_n)^3$ with respect to the ω-topology on S_n for all $n \in \mathbb{N}$, hence the same would occur for Φ and we are done. $\square$

Let $\mathcal{B}_S$ be a homology basis of $\mathcal{H}_1(S, \mathbb{Z})$ and label ν its cardinal number. Endow $\mathcal{F}_0^*(S)$ with the maximum norm, and consider the Fréchet differentiable map:

$$\mathcal{P} : \mathcal{F}_0^*(S) \times \mathcal{F}_0^*(S) \to \mathbb{C}^{3\nu}, \quad \mathcal{P}((h_1, h_2)) = \Big(\int_c ((e^{h_2 - h_1} - 1)\eta_1, (e^{h_2+h_1} - 1)\eta_2, (e^{h_2} - 1)\phi_3)\Big)_{c \in \mathcal{B}_S}.$$

The meromorphic data inside the integrals arise from multiplying g by e^{h_1} and ϕ_3 by e^{h_2}. Label $\mathcal{A}_0 : \mathcal{F}_0^*(S) \times \mathcal{F}_0^*(S) \to \mathbb{C}^{3\nu}$ as the Fréchet derivative of $\mathcal{P}$ at $(0, 0)$.

Claim 2.9. $\mathcal{A}_0|_{\mathcal{F}_0(W)\times\mathcal{F}_0(W)}$ *is surjective.*

Proof. Reason by contradiction and assume that $\mathcal{A}_0(\mathcal{F}_0(W) \times \mathcal{F}_0(W))$ lies in a complex subspace $\mathcal{U} = \{\big((x_c, y_c, z_c)\big)_{c\in\mathcal{B}_S} \in \mathbb{C}^{3\nu} \mid \sum_{c\in\mathcal{B}_S} \big(A_c x_c + B_c y_c + D_c z_c\big) = 0\}$, where A_c, B_c and $D_c \in \mathbb{C}$ for all $c \in \mathcal{B}_S$ and $\sum_{c\in\mathcal{B}_S} \big(|A_c| + |B_c| + |D_c|\big) \neq 0$. This simply means that:

$$-\int_{\Gamma_1} h\eta_1 + \int_{\Gamma_2} h\eta_2 = \int_{\Gamma_1} h\eta_1 + \int_{\Gamma_2} h\eta_2 + \int_{\Gamma_3} h\phi_3 = 0 \tag{2.1}$$

for all $h \in \mathcal{F}_0(W)$, where $\Gamma_1 = \sum_{c\in\mathcal{B}_S} A_c c$, $\Gamma_2 = \sum_{c\in\mathcal{B}_S} B_c c$ and $\Gamma_3 = \sum_{c\in\mathcal{B}_S} D_c c$.

Label $\Sigma_0 = \{f \in \mathcal{F}_0(W) \mid (f) \geq (\phi_3)^2\}$. By Theorem 2.3, the function $h = df/\phi_3 \in \mathcal{F}_0^*(S)$ lies in the closure of $\mathcal{F}_0(W)$ in the ω-topology on $\mathcal{F}_0^*(S)$ for any $f \in \Sigma_0$. Therefore, equation (2.1) can be applied formally to $h = df/\phi_3$, getting that $\int_{\Gamma_1} \frac{1}{g} df = \int_{\Gamma_2} g df = 0$ for all $f \in \Sigma_0$. Integrating by parts,

$$\int_{\Gamma_1} f \frac{dg}{g^2} = \int_{\Gamma_2} f\, dg = 0 \tag{2.2}$$

for all $f \in \Sigma_0$.

Let us show that $\Gamma_1 = 0$.

Let μ and b denote the genus of W and the number of ends of W. It is well known (see **[FK]**) that there exist $2\mu + b - 1$ cohomologically independent 1-forms in $\Omega_0(W)$ generating the first holomorphic De Rham cohomology group $\mathcal{H}^1_{\text{hol}}(W)$ of W. Thus, the map $H^1_{\text{hol}}(W) \longrightarrow \mathbb{C}^{2\mu+b-1}$, $\tau \mapsto \big(\int_c \tau\big)_{c\in B_0}$, where B_0 is any homology basis of W, is a linear isomorphism. Assume that $\Gamma_1 \neq 0$ and take $[\tau] \in H^1_{\text{hol}}(W)$ such that $\int_{\Gamma_1} \tau \neq 0$. Since W is an open surface, $\mathcal{F}_0(W)$ has infinite dimension and we can find $F \in \mathcal{F}_0(W)$ such that $(\tau + dF)_0 \geq (dg)_0 (g)^2_\infty (\phi_3)^2$. Set $h := \frac{(\tau+dF)g^2}{dg}$ and note that $(h) \geq (\phi_3)^2$. By Theorem 2.3, h lies in the closure of Σ_0 in $\mathcal{F}_0^*(S)$ with respect to the

ω-topology, hence equation (2.2) can be formally applied to h, giving that $\int_{\Gamma_1} \tau + dF = \int_{\Gamma_1} \tau = 0$, a contradiction.

By a similar argument $\Gamma_2 = 0$ and equation (2.1) becomes:

$$\int_{\Gamma_3} h\phi_3 = 0 \tag{2.3}$$

for all $h \in \mathcal{F}_0(W)$.

Since $\sum_{c\in\mathcal{B}_S} \big(|A_c| + |B_c| + |D_c|\big) \neq 0$, then $\Gamma_3 \neq 0$. Reason as above and choose $[\tau] \in H^1_{\rm hol}(W)$ and $F \in \mathcal{F}_0(W)$ such that $\int_{\Gamma_3} \tau \neq 0$ and $(\tau + dF)_0 \geq (\phi_3)$. Set $h := \frac{\tau+dF}{\phi_3}$ and note that $h \in \mathcal{F}_0^*(S)$. By Theorem 2.3, h lies in the closure of $\mathcal{F}_0(W)$ in $\mathcal{F}_0^*(S)$ with respect to the ω-topology, and equation (2.3) gives that $\int_{\Gamma_3} \tau + dF = \int_{\Gamma_3} \tau = 0$, a contradiction. This proves the claim. □

Let $\{e_1, \ldots, e_{3\nu}\}$ be a basis of $\mathbb{C}^{3\nu}$, fix $H_i = (h_{1,i}, h_{2,i}) \in \mathcal{A}_0^{-1}(e_i) \cap (\mathcal{F}_0(W) \times \mathcal{F}_0(W))$ for all i, and set $\mathcal{Q}_0 : \mathbb{C}^{3\nu} \to \mathbb{C}^{3\nu}$ as the analytical map given by

$$\mathcal{Q}_0((z_i)_{i=1,\ldots,3\nu}) = \mathcal{P}(\sum_{i=1,\ldots,3\nu} z_i H_i).$$

By Claim 2.9 $d(\mathcal{Q}_0)_0$ is an isomorphism, so there exists a closed Euclidean ball $U \subset \mathbb{C}^{3\nu}$ centered at the origin such that $\mathcal{Q}_0 : U \to \mathcal{Q}_0(U)$ is an analytical diffeomorphism. Furthermore, notice that $0 = \mathcal{Q}_0(0) \in \mathcal{Q}_0(U)$ is an interior point of $\mathcal{Q}_0(U)$.

On the other hand, by Lemmas 2.4 and 2.5 there exists a sequence $\{(f_n, \psi_n)\}_{n\in\mathbb{N}} \subset \mathcal{F}(W) \times \Omega_0(W)$ such that $(f_n) = (g)$ and $(\psi_n) = (\phi_3)$ for all n, and $\{(f_n, \psi_n)\}_{n\in\mathbb{N}} \to (g, \phi_3)$ in the ω-topology on $\mathcal{F}^*(S) \times \Omega_0^*(S)$.

Label $\mathcal{P}_n : \mathcal{F}_0^*(S) \times \mathcal{F}_0^*(S) \to \mathbb{C}^{3\nu}$ as the Fréchet differentiable map

$$\mathcal{P}_n((h_1, h_2)) = \Big(\int_c (e^{h_2-h_1}\eta_{1,n} - \eta_1, e^{h_2+h_1}\eta_{2,n} - \eta_2, e^{h_2}\psi_n - \phi_3)\Big)_{c\in\mathcal{B}_S},$$

where $\eta_{1,n} = \frac{1}{2}\psi_n(1/f_n - f_n)$ and $\eta_{2,n} = \frac{i}{2}\psi_n(1/f_n + f_n)$, and call $\mathcal{Q}_n : \mathbb{C}^{3\nu} \to \mathbb{C}^{3\nu}$ as the analytical map $\mathcal{Q}_n((z_i)_{i=1,\ldots,3\nu}) = \mathcal{P}_n(\sum_{i=1,\ldots,3\nu} z_i H_i)$ for all $n \in \mathbb{N}$. Since $\{\mathcal{Q}_n\}_{n\in\mathbb{N}} \to \mathcal{Q}_0$ uniformly on compacts subsets of $\mathbb{C}^{3\nu}$, without loss of generality we can suppose that $\mathcal{Q}_n : U \to \mathcal{Q}_n(U)$ is an analytical diffeomorphism and $0 \in \mathcal{Q}_n(U)$ for all n. Label $\mathbf{y_n} = (y_{1,n}, \ldots, y_{3\nu,n})$ as the unique point in U such that $\mathcal{Q}_n(\mathbf{y_n}) = 0$ and note that $\{\mathbf{y_n}\}_{n\in\mathbb{N}} \to 0$. Setting

$$g_n = e^{\sum_{j=1}^{3\nu} y_{j,n} h_{1,j}} f_n, \quad \phi_{3,n} = e^{\sum_{j=1}^{3\nu} y_{j,n} h_{2,j}} \psi_n$$

for all $n \in \mathbb{N}$, the sequence $\{(g_n, \phi_{3,n})\}_{n\in\mathbb{N}}$ solves the lemma. □

Corollary 2.10. *In the previous lemma we can choose $\phi_{3,n} = \phi_3$ for all $n \in \mathbb{N}$, provided that ϕ_3 extends holomorphically to W and ϕ_3 never vanishes on C_S.*

Proof. Without loss of generality, we can suppose that g, $1/g$ and dg never vanish on $\partial(M_S) \cup C_S$. Indeed, like in the proof of Claim 2.8 consider a sequence $\{M_n\}_{n\in\mathbb{N}}$ of tubular neighborhoods of M_S in W such that $\cap_{n=1}^{\infty} M_n = M_S$, $S_n := M_n \cup C_S$ is admissible in W, g, $1/g$ and dg never vanishes on $\partial(M_n)$ and $\gamma - M_n^\circ$ is a (non-empty) Jordan arc for any component γ of C_S, for all $n \in \mathbb{N}$. Let $h_n \in \mathcal{F}^*(S_n)$ be a smooth datum such that:

- $h_n|_{M_{S_n}} = g|_{M_{S_n}}$ and $\sum_{j=1}^{3} |\psi_{j,n}|^2$ never vanishes on S_n, where $\Psi_n = (\psi_{j,n})_{j=1,2,3} = \big(\frac{1}{2}(1/h_n - h_n), \frac{i}{2}(1/h_n + h_n), 1\big)\phi_3 \in \Omega_0^*(S_n)^3$, $n \in \mathbb{N}$,
- h_n, $1/h_n$ and dh_n never vanish on $\partial(M_{S_n}) \cup C_{S_n}$,
- $\Psi_n|_S - \Phi$ is exact on S, and
- the sequence $\{\Psi_n|_S\}_{n\in\mathbb{N}} \subset \Omega_0^*(S)^3$ converges to Φ in the ω-topology on S.

Reasoning as in the proof of Claim 2.8, if the Lemma held for any of the data in $\{\Psi_n \mid n \in \mathbb{N}\}$ the same would occur for Φ and we are done.

Reasoning like in the proof of Lemma 2.6, we can prove that $\hat{\mathcal{A}}_0 : \mathcal{F}_0(W) \to \mathbb{C}^\nu$ is surjective, where $\hat{\mathcal{A}}_0$ is the Fréchet derivative of $\hat{\mathcal{P}} : \mathcal{F}_0^*(S) \to \mathbb{C}^{2\nu}$, $\hat{\mathcal{P}}(h) := \mathcal{P}(h,0)$. Then take $\hat{H}_i \in \hat{\mathcal{A}}_0^{-1}(e_i) \cap \mathcal{F}_0(W)$ for all i, where $\{e_1, \ldots, e_{2\nu}\}$ is a basis of $\mathbb{C}^{2\nu}$, and define $\hat{\mathcal{Q}}_0 : \mathbb{C}^{2\nu} \to \mathbb{C}^{2\nu}$ by $\hat{\mathcal{Q}}_0((z_i)_{i=1,\ldots,2\nu}) = \hat{\mathcal{P}}(\sum_{i=1,\ldots,2\nu} z_i \hat{H}_i)$.

Use Riemann-Roch Theorem to find a holomorphic function $H \in \mathcal{F}_0(W)$ such that $(H) = (\phi_3|_{W-S})$, and then Lemma 2.4 to get $\{f_n\}_{n\in\mathbb{N}} \subset \mathcal{F}(W)$ such that $(f_n) = (g|_S)$ for all n and $\{f_n\}_{n\in\mathbb{N}} \to g/H$ in the ω-topology on $\mathcal{F}^*(S)$.

Set $\hat{\mathcal{P}}_n : \mathcal{F}_0^*(S) \to \mathbb{C}^{2\nu}$ by $\hat{\mathcal{P}}_n(h) = \big(\int_c (e^{-h}\eta_{1,n} - \eta_1, e^h \eta_{2,n} - \eta_2)\big)_{c\in\mathcal{B}_S}$, where $\eta_{1,n} = \frac{1}{2}\phi_3(\frac{1}{f_n H} - f_n H)$ and $\eta_{2,n} = \frac{i}{2}\phi_3(\frac{1}{f_n H} + f_n H)$, and call $\hat{\mathcal{Q}}_n : \mathbb{C}^{2\nu} \to \mathbb{C}^{2\nu}$ as the analytical map $\hat{\mathcal{Q}}_n((z_i)_{i=1,\ldots,2\nu}) = \hat{\mathcal{P}}_n(\sum_{i=1,\ldots,2\nu} z_i \hat{H}_i)$ for all $n \in \mathbb{N}$. To finish, reason as in the proof of Lemma 2.6. □

3. WEIERSTRASS REPRESENTATION AND FLUX MAP OF MINIMAL SURFACES

Let W be an open Riemann surface and let $X = (X_1, X_2, X_3) : W \to \mathbb{R}^3$ be a conformal minimal immersion. Denote by $\phi_j = \partial X_j$, $j = 1,2,3$, and $\Phi = \partial X \equiv (\phi_j)_{j=1,2,3}$. The 1-forms ϕ_k are holomorphic, have no real periods and satisfy that $\sum_{k=1}^3 \phi_k^2 = 0$. Furthermore, the intrinsic metric in W is given by $ds^2 = \sum_{k=1}^3 |\phi_k|^2$, hence ϕ_k, $k = 1,2,3$, have no common zeroes.

Conversely, a vectorial holomorphic 1-form $\Phi = (\phi_1, \phi_2, \phi_3)$ on W without real periods and satisfying that $\sum_{k=1}^3 \phi_k^2 = 0$ and $\sum_{k=1}^3 |\phi_k(P)|^2 \neq 0$ for all $P \in W$, determines a conformal minimal immersion $X : W \to \mathbb{R}^3$ by the expression:

$$X = \mathrm{Re} \int \Phi.$$

By definition, the triple Φ is said to be the Weierstrass representation of X. The meromorphic function $g = \frac{\phi_3}{\phi_1 - i\phi_2}$ corresponds to the Gauss map of X up to the stereographic projection (see **[Os]**).

We need the following definition:

Definition 3.1. Given a proper region $M \subset \mathcal{N}$ with possibly non empty boundary, we denote by $\mathcal{M}(M)$ the space of maps $X : M \to \mathbb{R}^3$ extending as a conformal minimal immersion to an open neighborhood of M in $\mathcal{N}$. This space will be equipped with the $\mathcal{C}^0$ topology of the uniform convergence on compact subsets of M.

Given $X \in \mathcal{M}(M)$ and an arclength parameterized curve $\gamma(s)$ in M, the conormal vector of X at $\gamma(s)$ is the unique unitary tangent vector $\mu(s)$ of X at $\gamma(s)$ such that $\{dX(\gamma'(s)), \mu(s)\}$ is a positive basis. If in addition γ is closed, the flux $p_X(\gamma)$ of X along γ is given by $\int_\gamma \mu(s)ds$, and it is easy to check that $p_X(\gamma) = \mathrm{Im} \int_\gamma \partial X$. The flux map $p_X : \mathcal{H}_1(M, \mathbb{Z}) \to \mathbb{R}^3$ is a group morphism.

Let $S \subset \mathcal{N}$ be a compact admissible subset. We denote by $\mathcal{J}(S)$ the space of smooth maps $X : S \to \mathbb{R}^3$ such that $X|_{M_S} \in \mathcal{M}(M_S)$ and $X|_{C_S}$ is regular, that is to say, $X|_\alpha$ is a regular curve for all $\alpha \subset C_S$. It is clear that $Y|_S \in \mathcal{J}(S)$ for all $Y \in \mathcal{M}(\mathcal{N})$.

Consider $X \in \mathcal{J}(S)$ and let $\varpi : C_S \to \mathbb{R}^3$ be an *smooth normal field* along C_S respect to X. This simply means that for any analytical arc $\alpha \subset C_S$ and any smooth parameter t on $X|_\alpha$, $\varpi(\alpha(t))$ is smooth, unitary and orthogonal to $(X|_\alpha)'(t)$, ϖ extends smoothly to any open analytical arc β in W containing α and ϖ is tangent to X on $\beta \cap S$. The normal field ϖ is said to be *orientable* respect to

X if for any component $\alpha \subset C_S$ having endpoints P_1, P_2 lying in $\partial(M_S)$, and for any arclength parameter s along $X|_\alpha$, the basis $B_i = \{(X|_\alpha)'(s_i), \varpi(s_i)\}$ of the tangent plane of $X|_{M_S}$ at P_i, $i = 1, 2$, are both positive or negative, where s_i is the value of s for which $\alpha(s_i) = P_i$, $i = 1, 2$.

Definition 3.2. We call $\mathcal{M}^*(S)$ as the space of marked immersions $X_\varpi := (X, \varpi)$, where $X \in \mathcal{J}(S)$ and ϖ is an orientable smooth normal field along C_S respect to X, endowed with the $\mathcal{C}^0$ topology of the uniform convergence of maps and normal fields.

Given $X_\varpi \in \mathcal{M}^*(S)$, let $\partial X_\varpi = (\hat{\phi}_j)_{j=1,2,3}$ be the complex vectorial "1-form" on S given by $\partial X_\varpi|_{M_S} = \partial(X|_{M_S})$, $\partial X_\varpi(\alpha'(s)) = dX(\alpha'(s)) + i\varpi(s)$, where α is a component of C_S and s is the arclength parameter of $X|_\alpha$ for which $\{dX(\alpha'(s_i)), \varpi(s_i)\}$ are positive, where as above s_1 and s_2 are the values of s for which $\alpha(s) \in \partial(M_S)$. If $(U, z = x + iy)$ is a conformal chart on N such that $\alpha \cap U = z^{-1}(\mathbb{R} \cap z(U))$, it is clear that $(\partial X_\varpi)|_{\alpha \cap U} = \big[dX(\alpha'(s)) + i\varpi(s)\big]s'(x)dz|_{\alpha \cap U}$, hence $\partial X_\varpi \in \Omega_0^*(S)^3$. Furthermore, $\hat{g} = \hat{\phi}_3/(\hat{\phi}_1 - i\hat{\phi}_2) \in \mathcal{F}^*(S)$ provided that $\hat{g}^{-1}(\infty) \subset S^\circ$.

It is clear that $\hat{\phi}_j$ is smooth on S, $j = 1, 2, 3$, and the same occurs for $\hat{g}$ on $S - \hat{g}^{-1}(\infty)$. Notice that $\sum_{j=1}^3 \hat{\phi}_j^2 = 0$, $\sum_{j=1}^3 |\hat{\phi}_j|^2$ never vanishes on S and $\text{Real}(\hat{\phi}_j)$ is an "exact" real 1-form on S, $j = 1, 2, 3$, hence we also have $X(P) = X(Q) + \text{Real} \int_Q^P (\hat{\phi}_j)_{j=1,2,3}$, P, $Q \in S$. For these reasons, $(\hat{g}, \hat{\phi}_3)$ will be called as the generalized "Weierstrass data" of X_ϖ. As $X|_{M_S} \in \mathcal{M}(M_S)$, then $(\phi_j)_{j=1,2,3} := (\hat{\phi}_j|_{M_S})_{j=1,2,3}$, and $g := \hat{g}|_{M_S}$ are the Weierstrass data and the meromorphic Gauss map of $X|_{M_S}$, respectively.

The group homomorphism

$$p_{X_\varpi} : \mathcal{H}_1(S, \mathbb{Z}) \to \mathbb{R}^3, \quad p_{X_\varpi}(\gamma) = \text{Im} \int_\gamma \partial X_\varpi,$$

is said to be the *generalized flux map* of X_ϖ. Obviously, $p_{X_{\varpi_Y}} = p_Y|_{\mathcal{H}_1(S)}$ provided that $X = Y|_S$ and ϖ_Y is the conormal field of Y along any curve in C_S.

4. PROPERNESS AND CONFORMAL STRUCTURE OF MINIMAL SURFACES

¿From now on, we label $x_k : \mathbb{R}^3 \to \mathbb{R}$ as the k-th coordinate function, $k = 1, 2, 3$. Given a compact $M \subset \mathcal{N}$ and a map $X : M \to \mathbb{R}^3$, we denote $\|X\| := \max_M \big\{ \big(\sum_{j=1}^3 (x_j \circ X)^2\big)^{1/2} \big\}$ as the maximum norm. For each ϱ, $a \in \mathbb{R}$, we call

$$\Pi_a(\varrho) = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_3 + \tan(\varrho)x_1 \leq a\}, \quad \Pi_a^*(\varrho) = \mathbb{R}^3 - \Pi_a(\varrho).$$

The following lemma concentrates most of the technical computations required in the proof of the main theorem of this section.

Lemma 4.1. *Let M, $V \subset \mathcal{N}$ be two compact regions with analytical boundary such that $M \subset V^\circ$ and $\mathcal{N} - M$ has no bounded components in $\mathcal{N}$.*

Consider $X \in \mathcal{M}(M)$ and let $p : \mathcal{H}_1(V, \mathbb{Z}) \to \mathbb{R}$ be any morphism extension of p_X. Suppose there are $\theta \in (0, \pi/4)$ and $\delta \in (0, +\infty)$ such that $X(\partial(M)) \subset \Pi_\delta^*(\theta) \cup \Pi_\delta^*(-\theta)$.

Then, for any $\epsilon > 0$ there exists $Y \in \mathcal{M}(V)$ such that $p_Y = p$, $\|Y - X\| < \epsilon$ on M, $Y(\partial(V)) \subset \Pi_{\delta+1}^*(\theta) \cup \Pi_{\delta+1}^*(-\theta)$ and $Y(V - M) \subset \Pi_\delta^*(\theta) \cup \Pi_\delta^*(-\theta)$.

Proof. We start with the following claim.

Claim 4.2. *The lemma holds when the Euler characteristic $\chi(V - M^\circ)$ vanishes.*

Proof. Since $M \subset V^\circ$ and $V^\circ - M$ has no bounded components in V°, then $V - M^\circ = \cup_{j=1}^k A_j$, where $A_1, \ldots, A_k$ are pairwise disjoint compact annuli. Write $\partial(A_j) = \alpha_j \cup \beta_j$, where $\alpha_j \subset \partial(M)$ and $\beta_j \subset \partial(V)$ for all j.

Since $X(\partial(M)) \subset \Pi_\delta^*(\theta) \cup \Pi_\delta^*(-\theta)$, each α_j can be divided into finitely many Jordan arcs α_j^i, $i = 1, \ldots n_j \geq 2$, laid end to end and satisfying that either $X(\alpha_j^i) \subset \Pi_\delta^*(\theta)$ or $X(\alpha_j^i) \subset \Pi_\delta^*(-\theta)$ for all i. Up to refining the partitions, we can assume that $n_j = m \in \mathbb{N}$ for all j.

An arc α_j^i is said to be *positive* if $X(\alpha_j^i) \subset \Pi_\delta^*(\theta)$, and *negative* otherwise. Notice that $X(\alpha_j^i) \subset \Pi_\delta^*(-\theta)$ for any negative (and possibly for some positive) α_j^i. We also label Q_j^i and Q_j^{i+1} as the endpoints of α_j^i, in such a way that $Q_j^{i+1} = \alpha_j^i \cap \alpha_j^{i+1}$, $i = 1, \ldots, m$, (obviously, $Q_j^{m+1} = Q_j^1$).

Let $\{r_j^i \mid i = 1, \ldots, m\}$ be a collection of pairwise disjoint analytical Jordan arcs in A_j such that r_j^i has initial point $Q_j^i \in \alpha_j$, final point $P_j^i \in \beta_j$, r_j^i is otherwise disjoint from $\partial(A_j)$, and r_j^i meets transversally α_j^i at Q_j^i, for all i and j.

Let W be a tubular neighborhood of V in $\mathcal{N}$, that is to say, an open connected subset of $\mathcal{N}$ such that $V \subset W$, $W - V$ consists of a finite collection of open annuli, and the closure of any annuli in $W - V$ intersects $\partial(V)$.

Set $M_0 = M \cup \left(\cup_{j,i} r_j^i \right)$, and notice that M_0 is an admissible subset in $\mathcal{N}$, and so in W. Call Ω_j^i as the closed disc in A_j bounded by $\alpha_j^i \cup r_j^i \cup r_j^{i+1}$ and a piece, named β_j^i, of β_j connecting P_j^i and P_j^{i+1}. Obviously $\Omega_j^i \cap \Omega_j^{i+1} = r_j^{i+1}$, $i < m$, $\Omega_j^m \cap \Omega_j^1 = r_j^1$, and $A_j = \cup_{i=1}^m \Omega_j^i$. The domain Ω_j^i is said to be positive (respectively, negative) if α_j^i is positive (respectively, negative).



FIGURE 1. The annulus A_j.

Take $\hat{X}_\varpi \in \mathcal{M}^*(M_0)$ such that $\hat{X}|_M = X$, and

$$\operatorname{dist}\left(\hat{X}(P_j^i)\,,\, \Pi_\delta(\theta) \cup \Pi_\delta(-\theta)\right) > 1, \tag{4.1}$$

for all i and j. In addition, we choose $\hat{X}_\varpi$ in such a way that:

(1_+) If Ω_j^i is positive then $\hat{X}(r_j^i \cup r_j^{i+1}) \subset \Pi_\delta^*(\theta)$.

(1_-) If Ω_j^i is negative then $\hat{X}(r_j^i \cup r_j^{i+1}) \subset \Pi_\delta^*(-\theta)$.

The existence of such a $\hat{X}_\varpi$ is standard, we omit the details.

We label $\hat{\phi}_k = \partial(x_k \circ \hat{X})$, $k = 1, 2, 3$, and recall that $\sum_{k=1}^3 \hat{\phi}_k^2 = 0$ and $\sum_{k=1}^3 |\hat{\phi}_k|^2$ never vanishes on M_0. Applying Lemma 2.6 to $\hat{\Phi} := (\hat{\phi}_k)_{k=1,2,3}$ on M_0 and integrating the resulting Weierstrass data on V, we can find $Y_0 \in \mathcal{M}(V)$ such that

$$p_{Y_0} = p_{\hat{X}_\varpi} = p_X = p \tag{4.2}$$

and

$$\|Y_0 - \hat{X}\| < \epsilon/3 \quad \text{on } M_0. \tag{4.3}$$

Furthermore, without loss of generality we can assume that

$$\operatorname{dist}\left(Y_0(P_j^i)\,,\, \Pi_\delta(\theta) \cup \Pi_\delta(-\theta)\right) > 1, \tag{4.4}$$

for all j, i, and

- (2_+) if Ω_j^i is positive then $Y_0(r_j^i \cup \alpha_j^i \cup r_j^{i+1}) \subset \Pi_\delta^*(\theta)$.
- (2_-) if Ω_j^i is negative then $Y_0(r_j^i \cup \alpha_j^i \cup r_j^{i+1}) \subset \Pi_\delta^*(-\theta)$.

This choice is possible thanks to the properties (4.1), (1_+) and (1_-).

Let $L^+ : \mathbb{R}^3 \to \mathbb{R}^3$ (respectively, $L^- : \mathbb{R}^3 \to \mathbb{R}^3$) denote the rotation of angle θ (respectively, $-\theta$) around the straight line parallel to the x_2-axis and containing the point $(0, 0, \delta)$. Note that $L^+(\Pi_\delta(\theta)) = \Pi_\delta(0)$, $L^+(\Pi_\delta(-\theta)) = \Pi_\delta(-2\theta)$, $L^-(\Pi_\delta(\theta)) = \Pi_\delta(2\theta)$ and $L^-(\Pi_\delta(-\theta)) = \Pi_\delta(0)$.

Call $Y^+ = L^+ \circ Y_0$ and $Y^- = L^- \circ Y_0 \in \mathcal{M}(V)$. Set $\mathcal{I}_+ = \{(i,j) \mid \Omega_j^i \text{ is positive}\}$ and $\mathcal{I}_- = \{(i,j) \mid \Omega_j^i \text{ is negative}\}$.

In a first step, we are going to deform the immersion Y^+. Roughly speaking, this deformation will be strong on $\cup_{(i,j)\in\mathcal{I}_+}\Omega_j^i$ and hardly modify Y^+ on $V - \cup_{(i,j)\in\mathcal{I}_+}\Omega_j^i$.

Label $\Psi = (\psi_k)_{k=1,2,3}$ as the Weierstrass data of Y^+ on V. For any $(i,j) \in \mathcal{I}_+$ let K_j^i and γ_j^i be a closed disc with analytical boundary and a compact analytical Jordan arc in Ω_j^i satisfying that:

(a1) $K_j^i \cap \partial(\Omega_j^i) \neq \emptyset$ and consists of an arc in $\beta_j^i - \{P_j^i, P_j^{i+1}\}$, $Y^+(\overline{\Omega_j^i - K_j^i}) \subset \Pi_\delta^*(0)$, and

$$\operatorname{dist}\left(Y^+(\beta_j^i - K_j^i)\,,\, \Pi_\delta(0) \cup \Pi_\delta(-2\theta)\right) > 1.$$

This choice is possible from (2_+) and (4.4) provided that K_j^i is chosen large enough in Ω_j^i.

(a2) The endpoints S_j^i and T_j^i of γ_j^i lie in $\alpha_j^i - \{Q_j^i, Q_j^{i+1}\}$ and $\partial(K_j^i) - \beta_j^i$, respectively, and γ_j^i is otherwise disjoint from $K_j^i \cup \partial(\Omega_j^i)$. Moreover, γ_j^i and α_j^i (resp., $\partial(K_j^i)$) meet transversally at S_j^i (resp., T_j^i) and ψ_3 never vanishes on γ_j^i. See Figure 1.

Set $S_+ = \left(\overline{V - \cup_{(i,j)\in\mathcal{I}_+}\Omega_j^i}\right) \cup \left(\cup_{(i,j)\in\mathcal{I}_+}(K_j^i \cup \gamma_j^i)\right)$, and notice that S_+ is admissible in $\mathcal{N}$, and so in W. Consider $\lambda^+ > 0$ such that

$$\operatorname{dist}\left((-\lambda^+, 0, 0) + Y^+(\cup_{(i,j)\in\mathcal{I}_+}K_j^i)\,,\, \Pi_\delta(-2\theta)\right) > 1. \tag{4.5}$$

This choice is possible because $\theta \in (0, \pi/4)$.

Claim 4.3. *There exist smooth data* $\hat{\Psi} = (\hat{\psi}_k)_{k=1,2,3} \in \Omega_0^*(S_+)^3$ *such that:*

(i) $\hat{\psi}_k = \psi_k$ *on* $\overline{S_+ - \cup_{(i,j)\in\mathcal{I}_+}\gamma_j^i}$ *for* $k = 1, 2$, *and* $\hat{\psi}_3 = (\psi_3)|_{S_+}$.

(ii) $\sum_{j=1}^3 (\hat{\psi}_k)^2 = 0$, *and* $\sum_{j=1}^3 |\hat{\psi}_k|^2$ *never vanishes on* S_+.

(iii) $Re\big(\int_{S_j^i}^{T_j^i}\hat{\psi}_1\big) = Re\big(\int_{S_j^i}^{T_j^i}\psi_1\big) - \lambda^+$, *where the integrals are computed along* γ_j^i, $(i,j)\in\mathcal{I}_+$.

Proof. For each $(i,j)\in\mathcal{I}_+$ consider an open conformal disc (U,z) on $\mathcal{N}$ (obviously depending on (i,j)) so that $\gamma_j^i\subset U$ and $z(\gamma_j^i)=[0,1]\subset\mathbb{R}$. Write $\psi_3(z)=f_3(z)dz$ on U and call g_+ as the meromorphic Gauss map of Y^+.

Let $\{\rho_t:\mathbb{R}\to\mathbb{C}\}_{t\in\mathbb{R}}$ be a smooth family of never vanishing smooth functions satisfying that:

- $\rho_t(x)=1$ if $x\in]-\infty,1/3]\cup[2/3,+\infty[$.
- If $|t|\geq 1$ then $\rho_t(x)=\frac{t}{g(x)f_3(x)}$ for all $x\in[\frac{1}{3}+\frac{1}{4+t^2},\frac{2}{3}-\frac{1}{4+t^2}]$.
- $A_0\leq|\rho_t(x)|\leq A_1|t|+A_2$ for all t, $x\in\mathbb{R}$, where A_0, A_1 and A_2 are suitable positive constants not depending on t.

Set $g_t=g\cdot(\rho_t\circ z)$ on γ_j^i for each $(i,j)\in\mathcal{I}_+$, and put $g_t=g_+$ on $\overline{S_+-\cup_{(i,j)\in\mathcal{I}_+}\gamma_j^i}$. Consider the generalized Weierstrass data $\hat{\Psi}_t=(\hat{\psi}_{k,t})_{k=1,2,3}\in\Omega_0^*(S_+)^3$ induced by the couple (g_t,ψ_3), and notice that $\hat{\Psi}_t$ satisfies (i) and (ii).

Since $\lim_{t\to+\infty}\mathrm{Re}\big(\int_{S_j^i}^{T_j^i}\hat{\psi}_{1,t}\big) = -\lim_{t\to-\infty}\mathrm{Re}\big(\int_{S_j^i}^{T_j^i}\hat{\psi}_{1,t}\big)=+\infty$, an intermediate value argument gives that $\mathrm{Re}\big(\int_{S_j^i}^{T_j^i}\hat{\psi}_{1,t_0}\big)=\mathrm{Re}\big(\int_{S_j^i}^{T_j^i}\psi_1\big)-\lambda^+$ for some $t_0\in\mathbb{R}$. It suffices to set $\hat{\Psi}=\hat{\Psi}_{t_0}$. ☐

Fix $P_0\in M$.

By Corollary 2.10, there exists $\{(\hat{\psi}_{1,n},\hat{\psi}_{2,n})\}_{n\in\mathbb{N}}\subset\Omega_0(V)^2$ converging to $(\hat{\psi}_1,\hat{\psi}_2)$ uniformly on S_+ and such that:

- $\sum_{k=1}^3(\hat{\psi}_{k,n})^2=0$ and $\sum_{k=1}^3|\hat{\psi}_{k,n}|^2$ never vanishes on V, where $\hat{\psi}_{3,n}=\hat{\psi}_3$, $\forall n\in\mathbb{N}$.
- The minimal immersion $Z_n^+\in\mathcal{M}(V)$ with initial condition $Z_n^+(P_0)=Y^+(P_0)$ and Weierstrass data $\hat{\Psi}_n=(\hat{\psi}_{k,n})_{k=1,2,3}$ is well defined and satisfies

$$p_{Z_n^+}=p_{Y^+},\quad \forall n\in\mathbb{N}. \tag{4.6}$$

Notice that $\{Z_n^+\}_{n\in\mathbb{N}}$ uniformly converges to Y^+ on $\overline{V-\cup_{(i,j)\in\mathcal{I}_+}\Omega_j^i}$, $\{x_1\circ Z_n^+\}_{n\in\mathbb{N}}$ uniformly converges to $x_1\circ Y^+-\lambda^+$ on $\cup_{(i,j)\in\mathcal{I}_+}K_j^i$, and $x_3\circ Z_n^+=x_3\circ Y^+$, $\forall n\in\mathbb{N}$ (see Claim 4.3). These facts let us find $n_0\in\mathbb{N}$ such that the following properties hold:

- $\|Z_{n_0}^+-Y^+\|<\epsilon/3$ on M.
- $Z_{n_0}^+\big(\cup_{(i,j)\in\mathcal{I}_+}\overline{\Omega_j^i-K_j^i}\big)\subset\Pi_\delta^*(0)$ and $Z_{n_0}^+\big(\cup_{(i,j)\in\mathcal{I}_+}(\beta_j^i-K_j^i)\big)\subset\Pi_{\delta+1}^*(0)$. See (a1).
- $\mathrm{dist}\big(Z_{n_0}^+(\cup_{(i,j)\in\mathcal{I}_+}K_j^i),\Pi_\delta(-2\theta)\big)>1$. Take into account (4.5).
- $\mathrm{dist}\big(Z_{n_0}^+(P_j^i),\Pi_\delta(-2\theta)\cup\Pi_\delta(0)\big)>1$, for any $(i,j)\in\mathcal{I}_-$. Use (4.4).
- $Z_{n_0}^+(r_j^i\cup\alpha_j^i\cup r_j^{i+1})\subset\Pi_\delta^*(-2\theta)$ for any $(i,j)\in\mathcal{I}_-$. It follows from (2_-).

Denote by $F^-:=L^-\circ L^-\circ Z_{n_0}^+\in\mathcal{M}(V)$. The above properties can be rewritten in terms of F^- as follows:

(A1) $\|F^--Y^-\|<\epsilon/3$ on M.
(A2) $F^-(\cup_{(i,j)\in\mathcal{I}_+}\overline{\Omega_j^i-K_j^i})\subset\Pi_\delta^*(2\theta)$ and $\mathrm{dist}\big(F^-(\cup_{(i,j)\in\mathcal{I}_+}(\beta_j^i-K_j^i)),\Pi_\delta(2\theta)\big)>1$.
(A3) $F^-\big(\cup_{(i,j)\in\mathcal{I}_+}K_j^i\big)\subset\Pi_{\delta+1}^*(0)$.
(A4) $\mathrm{dist}\big(F^-(P_j^i),\Pi_\delta(0)\cup\Pi_\delta(2\theta)\big)>1$, for any $(i,j)\in\mathcal{I}_-$.
(A5) $F^-(r_j^i\cup\alpha_j^i\cup r_j^{i+1})\subset\Pi_\delta^*(0)$, for any $(i,j)\in\mathcal{I}_-$.

Label $\Upsilon = (\varphi_k)_{k=1,2,3}$ as the Weierstrass data of F^- on V. Now, we are going to deform the immersion F^- in an analogous way. For any $(i,j) \in \mathcal{I}_-$ let K_j^i and γ_j^i be a closed disc and an analytical Jordan arc in Ω_j^i satisfying that:

(b1) $K_j^i \cap \partial(\Omega_j^i) \neq \emptyset$ and consists of a connected arc in β_j^i, $F^-(\overline{\Omega_j^i - K_j^i}) \subset \Pi_\delta^*(0)$, and
$$\operatorname{dist}\big(F^-(\beta_j^i - K_j^i)\,,\, \Pi_\delta(0) \cup \Pi_\delta(2\theta)\big) > 1.$$
This choice is possible from (A5) and (A4).

(b2) The endpoints S_j^i and T_j^i of γ_j^i lie in α_j^i and $\partial(K_j^i) - \beta_j^i$, respectively, and γ_j^i is otherwise disjoint from $K_j^i \cup \partial(\Omega_j^i)$. Moreover, γ_j^i and α_j^i (resp., $\partial(K_j^i)$) meet transversally at S_j^i (resp., T_j^i) and φ_3 never vanishes on γ_j^i.

Set $S_- = \big(\overline{V - \cup_{(i,j)\in\mathcal{I}_-}\Omega_j^i}\big) \cup \big(\cup_{(i,j)\in\mathcal{I}_-}(K_j^i \cup \gamma_j^i)\big)$. Consider $\lambda^- > 0$ such that
$$\operatorname{dist}\big((\lambda^-,0,0) + F^-(\cup_{(i,j)\in\mathcal{I}_-}K_j^i)\,,\, \Pi_\delta(2\theta)\big) > 1. \tag{4.7}$$

Reasoning as in Claim 4.3, there exist smooth data $\hat{\Upsilon} = (\hat{\varphi}_k)_{k=1,2,3} \in \Omega_0^*(S_-)^3$ satisfying that:

- $\hat{\varphi}_k = \varphi_k$ on $\overline{S_- - (\cup_{(i,j)\in\mathcal{I}_-}\gamma_j^i)}$, $k = 1,2$, and $\hat{\varphi}_3 = (\varphi_3)|_{S_-}$.
- $\sum_{k=1}^3 (\hat{\varphi}_k)^2 = 0$, and $\sum_{k=1}^3 |\hat{\varphi}_k|^2$ never vanishes on S_-.
- $\operatorname{Re}\big(\frac{1}{2}\int_{S_j^i}^{T_j^i} \hat{\varphi}_1\big) = \operatorname{Re}\big(\frac{1}{2}\int_{S_j^i}^{T_j^i} \varphi_1\big) + \lambda^-$, where as above we integrate along γ_j^i, $(i,j) \in \mathcal{I}_-$.

Again by Corollary 2.10, there exist $\{(\hat{\varphi}_{1,n}, \hat{\varphi}_{2,n})\}_{n\in\mathbb{N}} \subset \Omega_0(V)^2$ converging to $(\hat{\varphi}_1, \hat{\varphi}_2)$ uniformly on S_- and satisfying that:

- $\sum_{k=1}^3 (\hat{\varphi}_{k,n})^2 = 0$, and $\sum_{k=1}^3 |\hat{\varphi}_{k,n}|^2$ never vanishes on V, where $\hat{\varphi}_{3,n} = \hat{\varphi}_3$, $\forall n \in \mathbb{N}$.
- The immersion $G_n^- \in \mathcal{M}(V)$ with initial condition $G_n^-(P_0) = F^-(P_0)$ and Weierstrass data $\hat{\Upsilon}_n = (\hat{\varphi}_{k,n})_{k=1,2,3}$ is well defined and satisfies
$$p_{G_n^-} = p_{F^-}, \quad \forall n \in \mathbb{N}. \tag{4.8}$$

Arguing as above, we can find $n_1 \in \mathbb{N}$ such that the following properties hold:

(B1) $\|G_{n_1}^- - F^-\| < \epsilon/3$ on M.

(B2) $G_{n_1}^-\big(\cup_{(i,j)\in\mathcal{I}_-}\overline{\Omega_j^i - K_j^i}\big) \subset \Pi_\delta^*(0)$ and $G_{n_1}^-\big(\cup_{(i,j)\in\mathcal{I}_-}(\beta_j^i - K_j^i)\big) \subset \Pi_{\delta+1}^*(0)$. Use that $x_3 \circ G_{n_1}^- = x_3 \circ F^-$ on V and (b1).

(B3) $\operatorname{dist}\big(G_{n_1}^-(\cup_{(i,j)\in\mathcal{I}_-}K_j^i)\,,\, \Pi_\delta(2\theta)\big) > 1$. Take into account (4.7).

(B4) $G_{n_1}^-\big(\cup_{(i,j)\in\mathcal{I}_+}\overline{\Omega_j^i - K_j^i}\big) \subset \Pi_\delta^*(2\theta)$. See (A2).

(B5) $\operatorname{dist}\big(G_{n_1}^-(\cup_{(i,j)\in\mathcal{I}_+}(\beta_j^i - K_j^i))\,,\, \Pi_\delta(2\theta)\big) > 1$. It is implied by (A2).

(B6) $G_{n_1}^-\big(\cup_{(i,j)\in\mathcal{I}_+}K_j^i\big) \subset \Pi_{\delta+1}^*(0)$. It follows from (A3).

Summarizing, $\mathit{\Upsilon} := L^+ \circ G_{n_1} \in \mathcal{M}(V)$ satisfies the following properties:

- $\|\mathit{\Upsilon} - X\| \leq \|G_{n_1}^- - F^-\| + \|F^- - \mathit{\Upsilon}^-\| + \|\mathit{\Upsilon}_0 - \hat{X}\| < \epsilon$ on M. Use (A1), (B1) and (4.3).
- $\mathit{\Upsilon}\big(\cup_{(i,j)\in\mathcal{I}_-}\overline{\Omega_j^i - K_j^i}\big) \subset \Pi_\delta^*(-\theta)$ and $\mathit{\Upsilon}\big(\cup_{(i,j)\in\mathcal{I}_+}\overline{\Omega_j^i - K_j^i}\big) \subset \Pi_\delta^*(\theta)$. See (B2) and (B4).
- $\operatorname{dist}\big(\mathit{\Upsilon}(\cup_{(i,j)\in\mathcal{I}_-}(\beta_j^i - K_j^i))\,,\, \Pi_\delta(-\theta)\big) > 1$ and $\operatorname{dist}\big(\mathit{\Upsilon}(\cup_{(i,j)\in\mathcal{I}_+}(\beta_j^i - K_j^i))\,,\, \Pi_\delta(\theta)\big) > 1$. See (B2) and (B5).
- $\operatorname{dist}\big(\mathit{\Upsilon}(\cup_{(i,j)\in\mathcal{I}_-}K_j^i)\,,\, \Pi_\delta(\theta)\big) > 1$ and $\operatorname{dist}\big(\mathit{\Upsilon}(\cup_{(i,j)\in\mathcal{I}_+}K_j^i)\,,\, \Pi_\delta(-\theta)\big) > 1$. Use (B3) and (B6).

- $p_Y = p$. Take into account equations (4.2), (4.6) and (4.8).

This concludes the Claim. $\square$

To finish, we reason by induction on $s = -\chi(V - M^\circ) \in \mathbb{N} \cup \{0\}$. Claim 4.2 shows that the result holds for $s = 0$. Assume that the Lemma holds for $s = m$ and let us prove it for $s = m + 1$.

Since $s > 0$, there exists an analytic Jordan curve $\hat{\gamma} \in \mathcal{H}_1(V, \mathbb{Z}) - \mathcal{H}_1(M, \mathbb{Z})$ intersecting $V - M^\circ$ in a Jordan arc γ with endpoints $P_1, P_2 \in \partial(M)$ and otherwise disjoint from $\partial(M)$. Consider $F_\varpi \in \mathcal{M}^*(M \cup \gamma)$ such that $F|_M = X$, $F(\gamma) \subset \Pi^*_\delta(\theta) \cup \Pi^*_\delta(-\theta)$, and $p_{F_\varpi}(\hat{\gamma}) = p(\hat{\gamma})$. Here, we have taken into account that $F(P_i) \in \Pi^*_\delta(\theta) \cup \Pi^*_\delta(-\theta)$, for $i = 1, 2$. Notice that $p_{F_\varpi} = p|_{\mathcal{H}_1(M \cup \gamma, \mathbb{Z})}$. Label $\phi_k = \partial(x_k \circ F) \in \Omega^*_0(M \cup \gamma)$, $k = 1, 2, 3$.

Let W be a tubular neighborhood of V in $\mathcal{N}$. Applying Lemma 2.6 to $\Phi = (\phi_k)_{k=1,2,3}$, $S = M \cup \gamma$ and W, and integrating the resulting Weierstrass data, we can find a compact region M' with non empty analytical boundary and a minimal immersion $Z \in \mathcal{M}(M')$ such that $M \cup \gamma \subset (M')^\circ \subset M' \subset V^\circ$, $-\chi(V - (M')^\circ) = m$, $\mathcal{N} - M'$ has no bounded components in $\mathcal{N}$, $\|Z - X\| < \epsilon/2$ on M, $Z(\partial(M')) \subset \Pi^*_\delta(\theta) \cup \Pi^*_\delta(-\theta)$ and $p_Z = p|_{\mathcal{H}_1(M', \mathbb{Z})}$. Then, applying the induction hypothesis to M', V, Z, δ, and $\epsilon/2$, we obtain an immersion $Y \in \mathcal{M}(V)$ which satisfies the conclusion of the Lemma. $\square$

Theorem 4.4. *Let $p : \mathcal{H}_1(\mathcal{N}, \mathbb{Z}) \to \mathbb{R}^3$ and θ be a group morphism and a real number in $(0, \pi/4)$, respectively. Let $M \subset \mathcal{N}$ be a compact region of $\mathcal{N}$ such that $\mathcal{N} - M$ has no bounded components, and consider a non flat $Y \in \mathcal{M}(M)$ satisfying that $p_Y = p|_{\mathcal{H}_1(M, \mathbb{Z})}$ and $(x_3 + \tan(\theta)|x_1|) \circ Y > 1$.*

Then for any $\epsilon > 0$ there exists a conformal minimal immersion $X : \mathcal{N} \to \mathbb{R}^3$ satisfying the following properties:

- $p_X = p$,
- $(x_3 + \tan(\theta)|x_1|) \circ X : \mathcal{N} \to \mathbb{R}$ *is a positive proper function, and*
- $\|X - Y\| < \epsilon$ *on* M.

Proof. Without loss of generality, we will assume that $\epsilon < 1$.

Let $\{M_n \mid n \in \mathbb{N}\}$ be an exhaustion of $\mathcal{N}$ by compact regions with analytical boundary satisfying that $M_1 = M$, $M_n \subset M^\circ_{n+1}$ and $\mathcal{N} - M_n$ has no bounded components in $\mathcal{N}$ for all $n \in \mathbb{N}$.

Label $Y_1 = Y$, and by Lemma 4.1 and an inductive process, construct a sequence $\{Y_n\}_{n \in \mathbb{N}}$ of minimal immersions satisfying that

(a) $Y_n \in \mathcal{M}(M_n)$ for all $n \in \mathbb{N}$,
(b) $\|Y_n - Y_{n-1}\| < \epsilon/2^n$ on M_{n-1} for all $n \geq 2$,
(c) $p_{Y_n} = p|_{\mathcal{H}_1(M_n, \mathbb{Z})}$ for all $n \in \mathbb{N}$, and
(d) $Y_n(\partial(M_n)) \subset \Pi^*_n(\theta) \cup \Pi^*_n(-\theta)$ and $Y_n(M_n - M_{n-1}) \subset \Pi^*_{n-1}(\theta) \cup \Pi^*_{n-1}(-\theta)$ for all $n \geq 2$.

By items (a) and (b) and Harnack's theorem, $\{Y_n\}_{n \in \mathbb{N}}$ uniformly converges on compact subsets of $\mathcal{N}$ to a conformal minimal (possibly branched) immersion $X : \mathcal{N} \to \mathbb{R}^3$. Furthermore, $\|X - Y_n\| \leq \epsilon/2^n$ on M_n for all n. In particular from (d) we have that $(x_3 + \tan(\theta)|x_1|) \circ X \geq n - 1 - \epsilon/2^{n-1}$ on $M_n - M_{n-1}$, for all $n \geq 2$. On M_1 we also have that $(x_3 + \tan(\theta)|x_1|) \circ X \geq 1 - \epsilon > 0$, and so $(x_3 + \tan(\theta)|x_1|) \circ X$ is a positive proper function on $\mathcal{N}$.

Since the Weierstrass data of Y_n converge in the ω-topology to the ones of X, Hurwitz's theorem gives that either $X(\mathcal{N})$ is a point or X has no branch points. However, the condition $\|X - Y_1\| < \epsilon$ implies that $X(\mathcal{N})$ can not be a point provided that ϵ is small enough.

Finally, item (c) gives that $p_X = p$ and we are done. $\square$

5. PROPER MINIMAL SURFACES IN REGIONS WITH SUBLINEAR BOUNDARY

The main goal of this section is to prove the existence of proper hyperbolic minimal surfaces with non empty boundary in $\mathbb{R}^3$ contained in the region above a negative sublinear graph.

Throughout this section $\mathcal{N}$ will be the complex plane $\mathbb{C}$.

Theorem 5.1. *Let C denote the set $[-1,1]\times(0,1]\subset\mathbb{R}^2\equiv\mathbb{C}$ endowed with the conformal structure induced by $\mathbb{C}$.*[1]

Then there exists $X\in\mathcal{M}(C)$ satisfying that:

- $(x_1,x_3)\circ X : C\to\mathbb{R}^2$ *is proper.*
- $x_3\circ X > 0$ *on* $(x_1\circ X)^{-1}((-\infty,0])$, *and* $\lim_{n\to\infty} f(p_n) = 0$ *for any divergent sequence* $\{p_n\}_{n\in\mathbb{N}}$ *in C, where $f : C\to(-\infty,0]$ is the function* $f=\min\{\frac{x_3\circ X}{|x_1\circ X|+1},0\}$.

Proof. Let D_n denote the rectangle $[-2,2]\times[\frac{1}{n+1},2]\subset\mathbb{R}^2\equiv\mathbb{C}$, $n\in\mathbb{N}$. Label also $D=[-2,2]\times[0,2]$.

Lemma 5.2. *For any $\epsilon\in(0,1)$ there exists a sequence of non flat $X_k\in\mathcal{M}(D)$, $k\in\mathbb{N}$, such that*

(i) $\|(X_k-X_{k-1})\|<\epsilon/2^{k-1}$ *on* D_{k-1} *for all* $k\geq 2$.
(ii) $X_k([-2,2]\times\{\frac{1}{k+1}\})\subset\Pi_k^*(\frac{1}{k})$.
(iii) $X_k(D_k-D_{k-1})\subset\Pi_{k-1}^*(\frac{1}{k-1})\cup\Pi_k^*(\frac{1}{k})$ *for all* $k\geq 2$.
(iv) *If* $p\in D_k$ *and* $(x_1\circ X_k)(p)<0$, *then* $(x_3\circ X_k)(p)>1-\frac{\epsilon}{\sum_{j=1}^{k-1}2^j}>0$.

Proof. We are going to construct the sequence inductively. Take any non flat $X_1\in\mathcal{M}(D)$ satisfying that $X_1(D_1)\subset\Pi_1^*(1)$. Notice that Y_1 fulfills (ii) and (iv). Assume there exists a non flat immersion X_{n-1} satisfying (i), (ii), (iii) and (iv), and let us construct X_n.

Label $L:\mathbb{R}^3\to\mathbb{R}^3$ as the rotation of angle $\frac{1}{n-1}$ around the straight line parallel to the x_2-axis and containing the point $(0,0,n-1)$. Notice that $L(\Pi_{n-1}(\frac{1}{n-1}))=\Pi_{n-1}(0)$ and $L(\Pi_n(\frac{1}{n}))=\Pi_\zeta(-\frac{1}{n(n-1)})$, where $\zeta=n-1+\cos(1/n)\sec(1/(n^2-n))$.

Call $Y=L\circ X_{n-1}\in\mathcal{M}(D)$. From (ii), we have

$$Y([-2,2]\times\{1/n\})\subset\Pi_{n-1}^*(0). \tag{5.1}$$

By continuity and equation (5.1), there exists $\mu\in(\frac{1}{n+1},\frac{1}{n})$, close enough to $1/n$ so that

$$Y(\Theta)\subset\Pi_{n-1}^*(0),\quad\text{where }\Theta:=[-2,2]\times[\mu,1/n]. \tag{5.2}$$

Denote $\Delta:=[-2,2]\times[0,\mu]$. Notice that $D_n-D_{n-1}\subset\Delta\cup\Theta$, and $\varnothing=D_{n-1}\cap\Delta=D_{n-1}^\circ\cap\Theta^\circ=\Theta^\circ\cap\Delta^\circ$ (see figure 2).

Fix $\lambda>0$ such that

$$(-\lambda,0,0)+Y(\Delta)\subset\Pi_\zeta^*\Big(-\frac{1}{n(n-1)}\Big). \tag{5.3}$$

Denote $\Phi=(\phi_k)_{k=1,2,3}$ as the Weierstrass data of Y on D. Consider γ an analytic Jordan arc on Θ with endpoints $P\in\partial(D_{n-1})$ and $Q\in\partial(\Delta)$ and otherwise disjoint from $\partial(\Theta)$, and meeting transversally D_{n-1} and Δ. Moreover, we choose γ so that ϕ_3 never vanishes on γ. Label Λ as the admissible subset $D_{n-1}\cup\gamma\cup\Delta$ in $\mathbb{C}$ and consider $\hat\Phi=(\hat\phi_k)_{k=1,2,3}\in\Omega_0^*(\Lambda)^3$ satisfying

---

[1] By Caratheodory's Theorem, C is biholomorphic to the half disc $\overline{\mathbb{D}}_+$.



FIGURE 2. The sets in D

- $\hat{\phi}_k = \phi_k$ on $D_{n-1} \cup \Delta$, $\forall k = 1, 2$, and $\hat{\phi}_3 = \phi_3$ on D.
- $\sum_{k=1}^3 \hat{\phi}_k^2 = 0$ and $\sum_{k=1}^3 |\hat{\phi}_k|^2$ never vanishes.
- $\mathrm{Re}\big(\int_P^Q \hat{\phi}_1\big) = \mathrm{Re}\big(\int_P^Q \phi_1\big) - \lambda$, where the integrals are computed along γ.

The existence of $\hat{\Phi}$ follows by similar arguments to those used in Claim 4.3.

Let $W \subset \mathbb{C}$ be an open topological disc containing D, and without loos of generality, suppose that ϕ_3 extends holomorphically to W. We can apply Corollary 2.10 to the data W, $S = \Lambda$ and $\hat{\Phi} \in \Omega_0^*(\Lambda)^3$, to obtain a sequence $\{\hat{\Phi}_m = (\hat{\phi}_{k,m})_{k=1,2,3}\}_{m \in \mathbb{N}} \subset \Omega_0(D)^3$ converging to $\hat{\Phi}$ uniformly on Λ and such that $\hat{\phi}_{3,m} = \hat{\phi}_3 = \phi_3$, $\sum_{k=1}^3 \hat{\phi}_{k,m}^2 = 0$ and $\sum_{k=1}^3 |\hat{\phi}_{k,m}|^2$ never vanishes, for all $m \in \mathbb{N}$. Fix $P_0 \in D_{n-1}$, and consider the minimal immersion $Y_m \in \mathcal{M}(D)$ with initial condition $Y_m(P_0) = Y(P_0)$ and Weierstrass data $\hat{\Phi}_m$. Notice that $\{Y_m\}_{m \in \mathbb{N}}$ uniformly converges to Y on D_{n-1}, $\{x_1 \circ Y_m\}_{m \in \mathbb{N}}$ uniformly converges to $x_1 \circ Y - \lambda$ on Δ, and $x_3 \circ Y_m = x_3 \circ Y$, $\forall m \in \mathbb{N}$. As a consequence we can find $m_0 \in \mathbb{N}$ such that:

- $\|Y_{m_0} - Y\| < \epsilon/2^{n-1}$ on D_{n-1}.
- $Y_{m_0}(\Theta) \subset \Pi_{n-1}^*(0)$ (take into account (5.2)).
- $Y_{m_0}(\Delta) \subset \Pi_\zeta^*(-\frac{1}{n(n-1)})$. Use (5.3).

Define $X_n := L^{-1} \circ Y_{m_0} \in \mathcal{M}(D)$. From the above properties we get

(a) $\|X_n - X_{n-1}\| < \epsilon/2^{n-1}$ on D_{n-1}.
(b) $X_n(\Theta) \subset \Pi_{n-1}^*(\frac{1}{n-1})$.
(c) $X_n(\Delta) \subset \Pi_n^*(\frac{1}{n})$.

Property (a) directly gives (i). Since $[-2, 2] \times \{\frac{1}{n+1}\} \subset \Delta$, (c) implies (ii). Taking into account that $D_n - D_{n-1} \subset \Theta \cup \Delta$, (iii) follows from (b) and (c). Finally, property (iv) is an elementary consequence of (a), (b) and (c). □

From (i) and Harnack's theorem, the sequence $\{X_n\}_{n \in \mathbb{N}}$ uniformly converges on compact sets of $(-2, 2) \times (0, 2)$ to a conformal minimal (possibly branched) immersion $\hat{X} : (-2, 2) \times (0, 2) \to \mathbb{R}^3$. Since the Weierstrass data of X_n uniformly converge on compact sets of $(-2, 2) \times (0, 2)$ to the ones of $\hat{X}$, Hurwitz's theorem assures that either X has no branch points or $X((-2, 2) \times (0, 2))$ degenerates in a point. However, $\|\hat{X} - X_1\| < \epsilon$ on D_1, and so $\hat{X}$ does not degenerate provided that ϵ is small snough. Hence $\hat{X}$ is not branched and $X := \hat{X}|_C \in \mathcal{M}(C)$. Denote by $C_n =$

$[-1,1] \times [\frac{1}{n+1}, 1] \subset \mathbb{C}$, $n \in \mathbb{N}$. Using (i) and (iii) we get that $(x_3 + \tan(1)|x_1|) \circ X \geq n - 1 - 2\epsilon$ on $C_n - C_{n-1}$, for all $n \geq 2$. Therefore, $(x_1, x_3) \circ X : C \to \mathbb{R}^2$ is proper.

Consider $P \in C$ such that $(x_1 \circ X)(P) < 0$. For n large enough, $P \in C_n \subset D_n$ and $(x_1 \circ X_n)(P) < 0$ as well. Therefore (iv) gives $(x_3 \circ X)(P) \geq 1 - \epsilon > 0$ and $f(P) = 0$. Finally, consider a divergent sequence $\{P_n\}_{n \in \mathbb{N}}$ in C. For any $n \in \mathbb{N}$ we label $k(n) \in \mathbb{N}$ as the natural number such that $P_n \in C_{k(n)} - C_{k(n)-1}$ and note that $\{k(n)\}_{n \in \mathbb{N}}$ is divergent. From (i) and (iii), $(x_3 + \tan(\frac{1}{k(n)-1})(|x_1| + 1))(X(P_n)) > k(n) - 1 - 2\epsilon$. Hence, for n large enough,

$$0 \geq f(X(P_n)) \geq \min\left\{ \frac{k(n) - 1 - 2\epsilon}{|x_1(X(P_n))| + 1} - \tan\left(\frac{1}{k(n)-1}\right), 0 \right\} \geq -\tan\left(\frac{1}{k(n)-1}\right),$$

which converges to 0 as n goes to ∞.

The proof is done. $\square$

REFERENCES

[AS] L.V. Ahlfors and L. Sario, *Riemann surfaces.* Princeton Univ. Press, Princeton, New Jersey, 1960.

[Al] A. Alarcón, *On the existence of a proper conformal maximal disk in* $\mathbb{L}^3$. Differential Geom. Appl. **26** (2008), no. 2, 151-168.

[AFL] A. Alarcón, I. Fernández and F.J. López, *Complete minimal surfaces and harmonic functions.* Comment. Math. Helv. (In press).

[AFM] A. Alarcón, L. Ferrer and F. Martín, *Density theorems for complete minimal surfaces in* $\mathbb{R}^3$. Geom. Funct. Anal. **18 (1)** (2008), 1-49.

[Bi] E. Bishop, *Mappings of partially analytic spaces.* Amer. J. Math. **83** (1961), 209-242.

[CM] T.H. Colding and W.P. Minicozzi II. *The Calabi-Yau conjectures for embedded surfaces.* Ann. of Math. (2) **167** (2008), 211-243.

[CKMR] P. Collin, R. Kusner, W.H. Meeks III and H. Rosenberg, *The topology, geometry and conformal structures of properly embedded minimal surfaces* J. Differential Geom. **67** (2004), 377-393.

[DF] B. Drinovec Drnovšek and F. Forstnerič, *Holomorphic curves in complex spaces.* Duke Math. J. **139** (2007), 203-253.

[FK] H.M. Farkas and I. Kra, *Riemann surfaces.* Graduate Texts in Math. **71**, Springer Verlag, New York-Berlin, 1980.

[FMM] L. Ferrer, F. Martín and W.H. Meeks III, *Existence of proper minimal surfaces of arbitrary topological type.* Preprint (arXiv:0903.4194).

[HM] D. Hoffman and W.H. Meeks III, *The strong halfspace theorem for minimal surfaces.* Invent. Math. **101** (1990), 373-377.

[JM] L.P.M. Jorge and W.H. Meeks III, *The topology of complete minimal surfaces of finite total Gaussian curvature.* Topology **22** (1983), 203-221.

[JX] L.P.M. Jorge and F. Xavier, *A complete minimal surface in* $\mathbb{R}^3$ *between two parallel planes.* Ann. of Math. (2) **112** (1980), 203-206.

[Lo1] F.J. López, *Some Picard theorems for minimal surfaces.* Trans. Amer. Math. Soc. **356** (2004), 703-733.

[Lo2] F.J. López, *Uniform Approximation by algebraic minimal surfaces in* $\mathbb{R}^3$. Preprint (arXiv:0903.3209).

[LMM] F.J. López, F. Martín and S. Morales, *Adding handles to Nadirashvili's surfaces.* J. Differential Geom. **60** (2002), 155-175.

[LP] F.J. López and J. Pérez, *Parabolicity and Gauss map of minimal surfaces.* Indiana Univ. Math. J. **52** (2003), 1017-1026.

[MP1] W.H. Meeks III and J. Pérez, *Conformal properties in classical minimal surface theory.* Surveys in differential geometry. Vol. IX, 275–335, Surv. Differ. Geom., IX, Int. Press, Somerville, MA, 2004.

[MP2] W.H. Meeks III and J. Pérez, *Finite type annular ends for harmonic functions.* Preprint (arXiv:0909.1963).

[MPR] W.H. Meeks III, J. Pérez and A. Ros, *The embedded Calabi-Yau conjectures for finite genus.* In preparation.

[MUY] F. Martín, M. Umehara and K. Yamada, *Complete bounded null curves immersed in* $\mathbb{C}^3$ *and* $SL(2, \mathbb{C})$. Calc. Var. Partial Differential Equations **36** (2009), 119-139.

[Mo] S. Morales, *On the existence of a proper minimal surface in* $\mathbb{R}^3$ *with the conformal type of a disk.* Geom. Funct. Anal. **13** (2003), 1281-1301.

[Nad] N. Nadirashvili, *Hadamard's and Calabi-Yau's conjectures on negatively curved and minimal surfaces.* Invent. Math. **126** (1996), 457-465.

[Nar] R. Narasimhan, *Imbedding of holomorphically complete complex spaces.* Amer. J. Math. **82** (1960), 917-934.

[Ne] R.W. Neel, *Brownian motion and the parabolicity of minimal graphs.* Preprint (arXiv:0810.0669).

[Os] R. Osserman, *A survey of minimal surfaces.* Dover Publications, New York, second edition, 1986.

[Pe] J. Pérez, *Parabolicity and minimal surfaces.* Clay Math. Proc. **2**, Global theory of minimal surfaces, 163-174, Amer. Math. Soc., Providence, RI, 2005.

[Pi] G.P. Pirola, *Algebraic curves and non-rigid minimal surfaces in the Euclidean space.* Pacific J. Math. **183** (1998), 333-357.

[Re] R. Remmert, *Sur les espaces analytiques holomorphiquement séparables et holomorphiquement convexes.* C. R. Acad. Sci. Paris **243** (1956), 118-121.
[Ro] H.L. Royden, *Function Theory on Compact Riemann Surfaces.* J. Analyse Math. **18** (1967), 295-327.
[Sc1] S. Scheinberg, *Uniform approximation by functions analytic on a Riemann surface.* Ann. of Math. (2) **108** (1978), 257-298.
[Sc2] S. Scheinberg, *Uniform approximation by meromorphic functions having prescribed poles.* Math. Ann. **243** (1979), 83-93.
[SY] R. Schoen and S.T. Yau, *Lectures on Harmonic Maps.* Conference Proceedings and Lecture Notes in Geometry and Topology, II. International Press, Cambridge, MA, 1997.
[UY] M. Umehara and K. Yamada, *Maximal surfaces with singularities in Minkowski space.* Hokkaido Math. J. **35** (2006), 13-40.

DEPARTAMENTO DE MATEMÁTICA APLICADA, UNIVERSIDAD DE MURCIA, E-30100 ESPINARDO, MURCIA, SPAIN

E-mail address: `ant.alarcon@um.es`

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, UNIVERSIDAD DE GRANADA, E-18071 GRANADA, SPAIN

E-mail address: `fjlopez@ugr.es`